



1.0  
CONTACT HOUR

# Post hospitalization management of patients with COPD

By Daniel Blaha, MSN; Christine Vandenhouten, PhD, APHN-BC;  
Jolene Ramirez, MS, MSN, RNC-OB, NE-C; Julie Pomasl, MSN; and Hope Skarda, BSN, RN

One of the leading causes of morbidity and mortality in the United States, chronic obstructive pulmonary disease (COPD) impacts over 15 million Americans. In addition to lost lives, COPD has a significant economic and social burden.<sup>1,2</sup> It's estimated that the United States spends \$13.2 billion annually to manage patients with COPD, with hospital inpatient costs encompassing the majority of spending.<sup>3,4</sup> Managing COPD patients continues to challenge health systems. These challenges directly impact financial performance, with the Centers for Medicare and Medicaid Services imposing penalties on hospitals whose COPD readmission rates are in the bottom quartile.

Although multiple factors often unrelated to COPD contribute to readmission risk, health systems use many strategies to reduce the risk of readmission. Unfortunately, no clear data exist regarding how effective these strategies are at preventing read-

missions following a hospitalization for COPD.<sup>5,6</sup> Despite a lack of evidence supporting one approach to COPD readmission risk reduction, guidelines and recommendations exist regarding postacute interventions and services. This improvement project incorporated nursing interventions based on transitions theory and published guidelines from the American College of Chest Physicians and Canadian Thoracic Society into the design of a COPD readmission reduction initiative.<sup>7</sup>

## Setting

A 167-bed, Midwestern Level 3 trauma center was used for this improvement project. The pilot site is part of a larger multihospital system. Nationally, the readmission rate for COPD patients with Medicare is 20%.<sup>8</sup> The institution's COPD readmission rate in the 12-month period before the project was 22.6%, exceeding the target institutional COPD readmission



rate. To better understand factors influencing COPD readmission, an analysis of nursing-specific interventions, postacute follow-up care, and target medical interventions was completed. Results suggested that several gaps existed, including occasional lack of specialty follow-up, inconsistent prescribing of and adherence to scheduled and as-needed medications, and inconsistent postdischarge follow-up by the transition nurse.

Based on recent studies, it was noted that an opportunity existed

of a transitional management plan for patients with COPD discharged from the hospital setting. Professionals planning and supporting transitions must understand that transitions not only result from change, but also result in changes to health, relationships, and the individual's environment.<sup>14</sup> Because transition involves vulnerability and complex change in developmental, health, situational, and organizational areas, management of such transitions can't be addressed in an overly simplistic manner.<sup>14</sup>

the process of learning new skills, enhancing well-being, and improving confidence and coping during this critical period of transition.<sup>14,17-19</sup>

Transitions theory identifies important nursing therapeutics to support patient role transition, such as readiness assessment; preparation for transition, including provision of education; and role supplementation as patients transition.<sup>20</sup> Specific approaches using transitions theory to support patients with COPD include supporting health



***Patients with a COPD exacerbation being discharged home experience role transition requiring them to change not only their behavior, but also their definition of self.***

to reduce readmission rates among patients with COPD. These studies support the use of pulmonary rehabilitation; self-management education; specialty and primary care follow-up; medication management, particularly as it pertains to the use of inhalers; management of exacerbations; smoking cessation; and timely follow-up with a health coach or transition nurse to reduce COPD readmissions.<sup>1,2,7-12</sup>

### **Theoretical framework**

COPD has a complex, nonlinear disease trajectory, with varying and unpredictable patient adaptation to the disease process.<sup>13</sup> This project employed Meleis' transitions theory in the design

Psychosocial factors, as well as patient and health system complexity, were identified as causative factors for readmission.<sup>8,15</sup> That's why it's important for transition management to be individualized and holistic, including medical intervention, emotional support, and lifestyle promotion.<sup>13</sup>

Patients with a COPD exacerbation being discharged home experience role transition. Successfully moving through this transition requires patients to incorporate new knowledge and change not only their behavior, but also their definition of self.<sup>16</sup> By using transitions theory to guide patient interactions and care planning, nurses facilitate

state awareness, self-management, and ownership of symptom control; supporting the concept of hope; and ensuring social role fulfillment.<sup>17</sup> This approach utilizes the strengths of nursing therapeutic interventions in providing holistic management of patients as they move across the COPD continuum.

### **Project design**

This quality improvement project examined the impact of a transitions theory-based model by comparing pre- and postintervention readmission rates among COPD patients. The institution uses transitions-based care for chronic disease management. Transition nurses were already

in place and used to address the needs of patients with chronic conditions as they transitioned from inpatient to outpatient settings. Before the project, the transition nurse focused primarily on task-related checks, such as determining if patients obtained prescriptions, ensuring patients had a follow-up appointment, and checking to see if patients were experiencing exacerbation symptoms.

For this project, transition nurses received training on transitions theory, optimal COPD

As a member of the multidisciplinary team, the transition nurse worked with patients and healthcare providers to promote and facilitate optimal COPD management strategies. As part of the management plan, transition nurses completed follow-up phone calls 3 days' postdischarge, followed by weekly calls through 30 days' postdischarge. These calls used a structured assessment and interventions guided by previously identified nursing therapeutics. As compared with actions before the project, the

patient readmission rates. Patients who weren't initially identified as having a primary diagnosis of COPD but were later coded as such were also excluded because the transition nurse didn't have an opportunity to initiate all aspects of the project during the inpatient phase. To determine the impact of the use of a transition model on COPD readmission rates, results were compared with the readmission rate of COPD patients discharged home during the same time frame in 2016.



## ***Adherence to treatment regimens and medication concerns were correlated with readmission.***

management strategies, and targeted medical interventions based on guidelines from the American College of Chest Physicians and Canadian Thoracic Society. These guidelines outline elements required to optimize care of COPD patients postdischarge, including follow-up with a pulmonologist within 1 to 2 weeks of discharge, use of scheduled and as-needed medications, referral to a pulmonary rehabilitation clinic, use of home healthcare and telehealth, and creation of a self-management plan.<sup>7</sup> Self-management plans included exacerbation management, medication self-administration, healthcare provider follow-up, and smoking cessation.

transition nurse engaged the patient in discussions regarding self-management, including barriers to success and goal setting.

### **Data collection and analysis**

Institutional Review Board approval was received from the university and healthcare organization. Consenting patients discharged home with a primary diagnosis of COPD between January 12, 2017, and May 11, 2017, were included in this project. Patients discharged to a skilled nursing facility were neither included in the project nor in the comparative historical readmission data because the project examined the impact of transition nursing interventions on

The transition nurse tracked nursing-specific interventions, postacute medical follow-up care, and targeted medical interventions for each patient, noting whether the patient accepted or refused each. The transition nurse also recorded the readmission risk using the LACE Index Scoring Tool for Risk Assessment of Hospital Readmission, a validated tool that predicts the risk of readmission by calculating a risk score based on length of stay, acuity of admission, comorbidities, and use of ED services.<sup>21</sup> The transition nurse determined whether patients adhered to the self-management plan, defined as verbalizing the treatment plan; expressing no questions;



or use a medication, or stated intent not to follow the prescribed treatment plan. These data, along with demographic information, were analyzed to determine the impact each had on readmission rates.

The 30-day readmission rate was calculated for index COPD discharges occurring between January 12, 2017, and May 11, 2017, with comparison of readmission rates for the same time period in the previous year

existed between COPD patients who were readmitted and self-management plan nonadherence ( $p = 0.01$ ), expression of medication concerns ( $p = 0.02$ ), and a LACE score of 10 or greater ( $p = 0.04$ ).

#### Discussion

This quality improvement project identified several important findings. Although components of a COPD management program were in place, silos and coordina-

with COPD now consistently receive pulmonology consultation to ensure medication optimization. We've also seen a significant increase in the number of hospitalized COPD patients introduced to pulmonary rehabilitation through phase one completion. Unfortunately, the large majority of patients don't progress to phase two pulmonary rehabilitation due to lack of insurance coverage or patients not being sufficiently stable.



### ***Although all COPD patients are at risk for exacerbation and readmission, limited resources should target patients at the highest risk for readmission.***

(January 12, 2016, through May 11, 2016). This comparison eliminated seasonal factors that influence COPD exacerbation. Additional data included demographic information and whether the patient accepted or refused interventions ordered/offered.

SPSS 23.0 software was used to analyze factors influencing the COPD readmission rate 30 days' post hospital discharge.<sup>22</sup> Among patients discharged home, the readmission rate within the sample population was 15.2% ( $n = 46$ ) as compared with the 2016 readmission rate of 23.5% ( $n = 34$ ).

The Pearson's chi-squared test was used to determine the correlation between interventions and attributes and the readmission rate among patients discharged in 2017. A significant correlation

tion gaps between inpatient and outpatient services were discovered. The transition nurse could better coordinate inpatient and outpatient services, and support continuity in the care plan between pulmonary and hospitalist services. Use of a multidisciplinary readmission committee also supported effective implementation of proposed interventions and generated additional interventions for use with this patient population. For example, the committee identified the value of using online education modules for patient learning needs.

Gaps, including access to pulmonology, introduction to phase one pulmonary rehabilitation care, and use of effective educational tools, were identified and addressed. As a result of this project, patients

Use of transitions theory to guide nursing care was an effective approach to managing this complex disease. The framework guided a more structured process during in-person and phone interactions with patients, and ensured that conversations occurred around key transitional areas. The transition nurse is also continuing to develop education videos and materials for patients while in the hospital or at home.

It isn't surprising that adherence to treatment regimens and medication concerns were correlated with readmission. COPD is a complex, chronic disease that includes complex psychosocial factors. Nurses need to identify and develop nurse-sensitive interventions to assist patients, particularly those with a high

LACE score. Although all COPD patients are at risk for exacerbation and readmission, limited resources should target patients at the highest risk for readmission. Targeted interventions with this high-risk group have the greatest potential to reduce readmissions. For example, the transition nurse can provide additional

### Limitations

Results of this pilot project demonstrated factors impacting readmission of COPD patients; however, the small sample size and limited data collection period limit the generalizability of the findings. Although a reduction in the overall readmission rate was noted, comparison of a larger sample

transitions theory-based interventions supports optimized patient care throughout the continuum.

Nurse leaders need to ensure that transitions management is more than simply a list of tasks. Organizations benefit from a comprehensive transitions model, which includes structured processes and formalized roles that



***This quality improvement project demonstrated that evidence-based approaches to transition management of COPD patients can improve coordination of services and optimize care.***

psychosocial support to patients with a high LACE score.

This quality improvement project demonstrated that evidence-based approaches to transition management of COPD patients can improve coordination of services and optimize care. Additionally, use of a theoretical framework to guide the development of nursing interventions is important. Such a theoretical framework also takes greater advantage of nursing expertise based on the nursing process of assessment, care planning, intervention, and evaluation. It's essential to not only complete interactions with patients, but also engage them in self-management of their condition as emphasized through transitions theory. Ongoing development of this approach is warranted within the project setting and likely in many healthcare settings across the nation.

over a longer time span will provide further evidence to support the proposed interventions.

### Nursing implications

The utilization of transitions theory supported a more structured approach to care, resulting in improved nursing assessment, intervention, and patient participation. The use of transitions theory was effective in the ongoing evolution of transitions management processes and procedures. Additional education regarding transitions theory and related transitional care models will enhance the program and support patient and family engagement. It's also important to recognize that nursing care can't operate in isolation of other disciplines, including inpatient and outpatient healthcare providers. In fact, nurses are the key to the successful case management of chronic conditions. The use of

take advantage of nurses' skills, knowledge, and abilities.<sup>23</sup> It's incumbent on nurse leaders to ensure that new or current transitional care programs are evidence-based, continuum-focused, and grounded in nursing theory. **NM**

### REFERENCES

1. De Tratto K, Gomez C, Ryan CJ, Bracken N, Steffen A, Corbridge SJ. Nurses' knowledge of inhaler technique in the inpatient hospital setting. *Clin Nurse Spec.* 2014;28(3):156-160.
2. Parikh R, Shah TG, Tandon R. COPD exacerbation care bundle improves standard of care, length of stay, and readmission rates. *Int J Chron Obstruct Pulmon Dis.* 2016;11:577-583.
3. Khakban A, Sin DD, FitzGerald JM, et al. Ten-year trends in direct costs of COPD: a population-based study. *Chest.* 2015;148(3):640-646.
4. Shah T, Press VG, Huisinsh-Scheetz M, White SR. COPD readmissions: addressing COPD in the era of value-based health care. *Chest.* 2016;150(4):916-926.
5. Mannino DM, Thomashow B. Reducing COPD readmissions: great promise but big problems. *Chest.* 2015;147(5):1199-1201.

6. Hijawi SB, Abu Minshar M, Sharma G. Chronic obstructive pulmonary disease exacerbation: a single-center perspective on hospital readmissions. *Postgrad Med.* 2015;127(4):343-348.
7. Criner GJ, Bourbeau J, Diekemper RL, et al. Executive summary: prevention of acute exacerbation of COPD: American College of Chest Physicians and Canadian Thoracic Society Guideline. *Chest.* 2015;147(4):883-893.
8. Crisafulli E, Ortega S, Torres A. Predictors of readmission in a period of 30 days or less in acute exacerbation of chronic obstructive pulmonary disease. *Clin Pulm Med.* 2015;22(4):172-176.
9. Benzo R, Vickers K, Novotny PJ, et al. Health coaching and chronic obstructive pulmonary disease rehospitalization. A randomized study. *Am J Respir Crit Care Med.* 2016;194(6):672-680.
10. Cope K, Fowler L, Pogson Z. Developing a specialist-nurse-led 'COPD in-reach service'. *Br J Nurs.* 2015;24(8):441-445.
11. Johnson-Warrington V, Rees K, Gelder C, Morgan MD, Singh SJ. Can a supported self-management program for COPD upon hospital discharge reduce readmissions? A randomized controlled trial. *Int J Chron Obstruct Pulmon Dis.* 2016;11:1161-1169.
12. Raghavan D, Barter T, Joshi M. How to reduce hospital readmissions in chronic obstructive pulmonary disease? *Curr Opin Pulm Med.* 2016;22(2):106-112.
13. Cornforth A. COPD self-management supportive care: chaos and complexity theory. *Br J Nurs.* 2013;22(19):1101-1104.
14. Meleis AI, Sawyer LM, Im EO, Hilfinger Messias DK, Schumacher K. Experiencing transitions: an emerging middle-range theory. *ANS Adv Nurs Sci.* 2000;23(1):12-28.
15. Coventry PA, Gemmell I, Todd CJ. Psychosocial risk factors for hospital readmission in COPD patients on early discharge services: a cohort study. *BMC Pulm Med.* 2011;11:49.
16. Meleis AI. Role insufficiency and role supplementation: a conceptual framework. In: Meleis AI, ed. *Transitions Theory: Middle-Range and Situation-Specific Theories in Nursing Research and Practice.* New York, NY: Springer Publishing Company; 2010:13-24.
17. Geary CR, Schumacher KL. Care transitions: integrating transition theory and complexity science concepts. *ANS Adv Nurs Sci.* 2012;35(3):236-248.
18. Halting AG, Heggdal K. Patients' experiences of health transitions in pulmonary rehabilitation. *Nurs Inq.* 2012;19(4):345-356.
19. Im EO. Situation-specific theories from the middle-range transitions theory. *ANS Adv Nurs Sci.* 2014;37(1):19-31.
20. Schumacher KL, Meleis AI. Transitions: a central concept in nursing. In: Meleis AI, ed. *Transitions Theory: Middle-Range and Situation-Specific Theories in Nursing Research and Practice.* New York, NY: Springer Publishing Company; 2010:38-51.
21. van Walraven C, Dhalla IA, Bell C, et al. Derivation and validation of an index to predict early death or unplanned readmission after discharge from hospital to the community. *CMAJ.* 2010;182(6):551-557.
22. IBM Corp. *IBM SPSS Statistics for Windows, Version 23.0.* Armonk, NY: IBM Corp.; 2015.
23. Dusek B, Pearce N, Harripaul A, Lloyd M. Care transitions: a systematic review of best practices. *J Nurs Care Qual.* 2015;30(3):233-239.

In Green Bay, Wisc., Daniel Blaha is a nurse leader in hospice at Aurora at Home; Christine Vandenhouten is the chair/associate professor of nursing and health studies at the University of Wisconsin—Green Bay; and Jolene Ramirez is a director of inpatient nursing. Julie Pomasi is the director of quality management, and Hope Skarda is a transition of care RN at Aurora BayCare Medical Center.

The authors and planners have disclosed no potential conflicts of interest, financial or otherwise.

DOI-10.1097/01.NUMA.0000527713.06404.d1

For more than 146 additional continuing education articles related to management topics, go to [NursingCenter.com/CE](http://NursingCenter.com/CE).

**CE CONNECTION**

**Earn CE credit online:**  
Go to <http://nursing.ceconnection.com> and receive a certificate *within minutes*.

#### INSTRUCTIONS

### Post hospitalization management of patients with COPD

#### TEST INSTRUCTIONS

- Read the article. The test for this CE activity is to be taken online at <http://nursing.ceconnection.com>.
- You'll need to create (it's free!) and login to your personal CE Planner account before taking online tests. Your planner will keep track of all your Lippincott Professional Development online CE activities for you.
- There's only one correct answer for each question. A passing score for this test is 13 correct answers. If you pass, you can print your certificate of earned contact hours and access the answer key. If you fail, you have the option of taking the test again at no additional cost.
- For questions, contact Lippincott Professional Development: 1-800-787-8985.
- Registration deadline is **January 31, 2020**.

#### PROVIDER ACCREDITATION

Lippincott Professional Development will award 1.0 contact hour for this continuing nursing education activity.

Lippincott Professional Development is accredited as a provider of continuing nursing education by the American Nurses Credentialing Center's Commission on Accreditation.

This activity is also provider approved by the California Board of Registered Nursing, Provider Number CEP 11749 for 1.0 contact hour, and the District of Columbia, Georgia, and Florida CE Broker #50-1223.

Payment: The registration fee for this test is \$12.95.